

## Commentary on: Are we overpathologizing everyday life? A tenable blueprint for behavioral addiction research

### *On the slippery slopes: The case of gambling addiction*

LUKE CLARK\*

Centre for Gambling Research at UBC, Department of Psychology, University of British Columbia, Vancouver, Canada

(Received: April 24, 2015; revision received: April 28, 2015; accepted: April 30, 2015)

---

Billieux et al. (2015) propose that the recent proliferation of behavioral addictions has been driven by deficiencies in the underlying research strategy. This commentary considers how pathological gambling (now termed gambling disorder) traversed these challenges to become the first recognized behavioral addiction in the DSM-5. Ironically, many similar issues continue to exist in research on gambling disorder, including question-marks over the validity of tolerance, heterogeneity in gambling motives, and the under-specification of neuroimaging biomarkers. Nevertheless, I contend that the case for gambling disorder as a behavioral addiction has been bolstered by the existence of clear and consistent functional impairment (primarily in the form of debt), coupled with the development of a public health approach that has given emphasis to product features (i.e. the structural characteristics of gambling forms) as much as individual dispositions (the ‘addictive personality’).

**Keywords:** pathological gambling, video games, addiction, tolerance, neuroimaging, structural features

---

Billieux, Schimmenti, Khazaal, Maurage and Heeren (2015) articulately explain how the recent proliferation of soft behavioral addictions could arise from a circular research approach, involving 3 stages. The first step is the derivation of a screening tool, based on adapting established criteria for substance use disorders. The second step is the confirmation in epidemiological datasets that some prevalence of the putative addiction exists in the general population. The third step is the testing for neurocognitive markers in the putative addiction, when those markers are themselves derived from the same essential criteria (albeit in substance use disorders). The widespread emergence of this approach has likely been fueled by the ratification of ‘behavioral addiction’ via the case of gambling disorder (previously termed pathological gambling) in the DSM-5 (Petry et al., 2014).

Billieux et al. (2015) imply that this current conceptualization of gambling disorder was validated by “decades of empirical research”. Reading their article, I was prompted to reconsider how (or indeed whether) gambling disorder successfully overcame these hurdles that other putative behavioral addictions now face. Certainly, similar controversy existed around the time that the DSM-III introduced the pathological gambling diagnosis in 1980 (Lesieur, 1984). Indeed, when taken in isolation, many specific lines of evidence in gambling disorder remain open to criticisms analogous to those raised by Billieux et al. (2015). I will consider three examples. First, Billieux et al. (2015) question the validity of tolerance in behavioral addictions. Tolerance is often regarded as one of the hallmarks of an addiction syndrome (Shaffer et al., 2004). In the case of gambling, individuals with gambling disorder clearly play with progressively larger sums of money over time, and many screening instruments include an item that refers to escalating bet size. However, this effect may have a distinct motivation

from the phenomenon in substance use disorders, where an opponent process causes the addicted individual to require higher doses of drug to achieve the same subjective effect. In gambling disorder, it is not clear that the increasing wagers are necessary for need satisfaction; an accumulation of debt may entirely justify the need to escalate one’s bet (Blaszczynski, Walker, Sharpe & Nower, 2008).

Second, Billieux et al. (2015) propose that a common symptom presentation (e.g. of multiplayer online game addiction) can arise from profoundly different psychological motives. Such heterogeneity in motives clearly also exists in gambling disorder. Factor analytic studies differentiate coping motives (gambling to escape), enhancement motives (gambling for excitement) and socializing motives (Stewart & Zack, 2008), with some dispute over the place of money as a motivating factor (Flack & Morris, 2014). Different preferred forms of gambling and mental health co-morbidities may align with these motives (Blaszczynski & Nower, 2002) with implications for treatment tailoring.

Third, the search for biomarkers for gambling disorder has entered an interesting phase. Using functional MRI, a substantial number of studies have focused attention on the ‘brain reward system’ (centered on the ventral striatum and medial prefrontal cortex) but the direction of group differences in these regions is inconsistent (Limbrick-Oldfield, Van Holst & Clark, 2013). A similar number of studies describe over- and under-activity in groups with gambling disorder; results tend to be interpreted with two distinct

---

\* Corresponding address: Dr Luke Clark; Centre for Gambling Research at UBC, Department of Psychology, University of British Columbia, 2136 West Mall, Vancouver, BC, Canada V6T 1Z4; E-mail: luke.clark@psych.ubc.ca

theoretical positions, reward deficiency versus incentive salience (Leyton & Vezina, 2013). PET data looking at brain neurochemistry are even stranger, where dopamine tracers yield little overlap in the signature of gambling disorder and substance use disorders (Boileau et al., 2014; Clark et al., 2012). One possibility is that the effects in substance use disorders are drug-induced (e.g. cocaine neurotoxicity). Another distinct possibility is that the biomedical model of addictions represent an over-simplification in the case of gambling disorder (c.f. Hall, Carter & Forlini, 2015; Nutt, Lingford-Hughes, Erritzoe & Stokes, 2015).

One salient riposte to these debates comes in the form of functional impairment. For clinicians, this point goes without saying; clients do not present to services seeking help without some form of functional impairment (this may be perceived by family or the legal system rather than directly by the individual). In the case of gambling disorder, financial debt is ubiquitous, crippling, and pernicious, drawing friends and family quickly into the spiral of harms. It is not uncommon for individuals with gambling disorder to become homeless, attempt suicide or resort to crime, as direct responses to their mounting debts (Clark & Walker, 2009; Manning et al., 2015; Sharman, Dreyer, Aitken, Clark & Bowden-Jones, 2015). These objective harms contrast with the subjective, transient and contextual distress that is often described for other putative behavioral addictions.

A public health model provides a further perspective, by emphasizing how harms arise through the interaction of individual risk factors and features of the addictive agent (Korn & Shaffer, 1999). In the specific case of gambling, this can be expressed as the interplay of the ‘gambler and the game’, or the ‘player and the product’ (Clark, 2014). The addictions framework is grounded in a biomedical model that may give disproportionate weight to individual vulnerabilities, which effectively aims to characterize the ‘addictive personality’ in neurobiological and psychological terms. But within this framework – and in the continued absence of clear biomarkers – any excessive behavior can be considered an addiction, and thus the slippery slope beckons. By understanding how product features act on the consumer, this may help impose some thresholds on the slippery slope. In the case of drug addiction, it is clear for example that the addictive potential of tobacco is vastly different when nicotine is administered in the form of cigarettes as opposed to chewing tobacco (Edwards, 2005). In the case of gambling games, various structural characteristics have been identified, such as speed of play (Chóliz, 2010), near-miss features (Clark, Lawrence, Astley-Jones & Gray, 2009) or illusion of control devices (Ladouceur & Sevigny, 2005). There is increasing evidence that at least some of these features are linked to the addictive potential of different forms of gambling, and modulate brain activity in the same regions affected by individual vulnerability factors.

A similar approach is proving fruitful in the case of video gaming (especially online gaming). Some features are shared across gambling games and video games (e.g. reinforcement schedules), while others are specific (e.g. advancement of the avatar in video games) (King, Delfabbro & Griffiths, 2009). In the case of ‘food addiction’ or binge eating, it may also be possible to relate sugar, salt or fat content of foods to behavioral models of consumption in a way that at least generates a tractable program of research

and falsifiable hypotheses (Avena, Gearhardt, Gold, Wang & Potenza, 2012; Ziauddeen, Farooqi & Fletcher, 2012). It is less convincing whether or how such product features should be conceptualized for other putative behavioral addictions, such as ‘work addiction’ or ‘tango addiction’. Thus, and in conclusion, any blueprint for behavioral addictions research would benefit from an increased emphasis on the psychological properties of addictive products.

---

*Funding sources:* The Centre for Gambling Research at UBC is supported by funding from the Province of British Columbia and the British Columbia Lottery Corporation (BCLC). Dr Clark holds research funding in the UK from the Medical Research Council (G1100554).

*Conflict of interest:* Dr Clark is the Director of the Centre for Gambling Research at UBC, which is supported by funding from the Province of British Columbia and the British Columbia Lottery Corporation (BCLC). This support imposes no constraints on publishing. Dr Clark has provided paid consultancy to Cambridge Cognition Ltd. on issues relating to neurocognitive assessment. Dr Clark has not received any other direct or indirect payments from the gambling industry or any other groups substantially funded by gambling to conduct research or to speak at conferences or events.

---

## REFERENCES

- Avena, N. M., Gearhardt, A. N., Gold, M. S., Wang, G.-J. & Potenza, M. N. (2012). Tossing the baby out with the bathwater after a brief rinse? The potential downside of dismissing food addiction based on limited data. *Nature Reviews Neuroscience*, *13*(7), 514–514.
- Billieux, J., Schimmenti, A., Khazaal, Y., Maurage, P. & Heeren, A. (2015). Are we overpathologizing everyday life? A tenable blueprint for behavioral addiction research. *Journal of Behavioural Addictions*, *4*, 119–123.
- Blaszczynski, A. & Nower, L. (2002). A pathways model of problem and pathological gambling. *Addiction*, *97*(5), 487–499.
- Blaszczynski, A., Walker, M., Sharpe, L. & Nower, L. (2008). Withdrawal and tolerance phenomenon in problem gambling. *International Gambling Studies*, *8*(2), 179–192.
- Boileau, I., Payer, D., Chugani, B., Lobo, D. S. S., Houle, S., Wilson, A. A., Warsh, J., Kish, S. J. & Zack, M. (2014). In vivo evidence for greater amphetamine-induced dopamine release in pathological gambling: A positron emission tomography study with [(11)C]-(+)-PHNO. *Molecular Psychiatry*, *19*, 1305–1313.
- Chóliz, M. (2010). Experimental analysis of the game in pathological gamblers: Effect of the immediacy of the reward in slot machines. *Journal of Gambling Studies*, *26*(2), 249–256.
- Clark, L. (2014). Disordered gambling: The evolving concept of behavioral addiction. *Annals of the New York Academy of Sciences*, *1327*(1), 46–61.
- Clark, L., Lawrence, A. J., Astley-Jones, F. & Gray, N. (2009). Gambling near-misses enhance motivation to gamble and recruit win-related brain circuitry. *Neuron*, *61*(3), 481–490.
- Clark, L., Stokes, P. R., Wu, K., Michalczuk, R., Benecke, A., Watson, B. J., Egerton, A., Piccini, P., Nutt, D. J., Bowden-Jones, H.

- & Lingford-Hughes, A. R. (2012). Striatal dopamine D(2)/D(3) receptor binding in pathological gambling is correlated with mood-related impulsivity. *NeuroImage*, *63*(1), 40–46.
- Clark, C. & Walker, D. M. (2009). Are gamblers more likely to commit crimes? An empirical analysis of a nationally representative survey of US young adults. *International Gambling Studies*, *9*(2), 119–134.
- Edwards, G. (2005). *Matters of substance: Drugs—and why everyone's a user*. London, U.K: Macmillan.
- Flack, M. & Morris, M. (2014). Problem gambling: One for the money...? *Journal of Gambling Studies*. doi:10.1007/s10899-014-9484-z
- Hall, W., Carter, A. & Forlini, C. (2015). The brain disease model of addiction: Is it supported by the evidence and has it delivered on its promises? *The Lancet Psychiatry*, *2*(1), 105–110.
- King, D., Delfabbro, P. & Griffiths, M. (2009). Video Game structural characteristics: A new psychological taxonomy. *International Journal of Mental Health and Addiction*, *8*(1), 90–106.
- Korn, D. A. & Shaffer, H. J. (1999). Gambling and the health of the public: Adopting a public health perspective. *Journal of Gambling Studies*, *15*(4), 289–365.
- Ladouceur, R. & Sevigny, S. (2005). Structural characteristics of video lotteries: Effects of a stopping device on illusion of control and gambling persistence. *Journal of Gambling Studies*, *21*(2), 117–131.
- Lesieur, H. R. (1984). *The chase: The compulsive gambler*. Rochester, VT: Schenkman.
- Leyton, M. & Vezina, P. (2013). Striatal ups and downs: Their roles in vulnerability to addictions in humans. *Neuroscience and Biobehavioral Reviews*. doi: 10.1016/j.neubiorev.2013.01.018
- Limbrick-Oldfield, E. H., Van Holst, R. J. & Clark, L. (2013). Fronto-striatal dysregulation in drug addiction and pathological gambling: Consistent inconsistencies? *NeuroImage: Clinical*, *2*, 385–393.
- Manning, V., Kee, P., Yang, Y., Ng, A., Guo, S., Kandasami, G. & Eng, K. (2015). Suicidal ideation and lifetime attempts in substance and gambling disorders. *Psychiatry Research*, *225*(3), 706–709.
- Nutt, D. J., Lingford-Hughes, A., Erritzoe, D. & Stokes, P. R. A. (2015). The dopamine theory of addiction: 40 years of highs and lows. *Nature Reviews Neuroscience*, *16*(5), 305–312.
- Petry, N. M., Blanco, C., Auriacombe, M., Borges, G., Buchholz, K., Crowley, T. J. Grant, B. F., Hasin, D. S. & O'Brien, C. (2014). An overview of and rationale for changes proposed for pathological gambling in DSM-5. *Journal of Gambling Studies*, *30*(2), 493–502.
- Shaffer, H. J., LaPlante, D. A., LaBrie, R. A., Kidman, R. C., Donato, A. N. & Stanton, M. V. (2004). Toward a syndrome model of addiction: Multiple expressions, common etiology. *Harvard Review of Psychiatry*, *12*(6), 367–374.
- Sharman, S., Dreyer, J., Aitken, M., Clark, L. & Bowden-Jones, H. (2015). Rates of problematic gambling in a British homeless sample: A preliminary study. *Journal of Gambling Studies*, *31*(2), 525–532.
- Stewart, S. H. & Zack, M. (2008). Development and psychometric evaluation of a three-dimensional Gambling Motives Questionnaire. *Addiction*, *103*(7), 1110–1117.
- Ziauddeen, H., Farooqi, I. S. & Fletcher, P. C. (2012). Obesity and the brain: How convincing is the addiction model? *Nature Reviews Neuroscience*, *13*(4), 279–286.